

For wound management

ALLEVYN[◊] LIFE Foam Dressings are intended for exuding wounds including:

- pressure injuries
- diabetic foot ulcers
- first- and second-degree burns
- fungating ulcers
- leg ulcers
- surgical wounds
- donor sites
- skin tears

ALLEVYN LIFE may be used on infected wounds, which should be treated per local clinical protocol. The sacrum shape suits the sacrum/coccyx and can be worn for up to five days.^{3,4}

Advantages

- Maintains an optimal moist wound environment.²
- May help reduce the risk of maceration.⁶
- Reduced dressing change visits by 50% in a study.^{6*}

For pressure injury prevention

ALLEVYN LIFE offers a five-layer foam design that helps protect against pressure injuries as part of a pressure injury protocol.⁷

Advantages

- Reduced incidence of sacral pressure injuries by 71% compared to standard preventive care alone, in an independent randomized controlled trial.^{7,8}
- Showed a cost saving of 69% per patient compared to using standard preventive care alone, in an associated economic study.⁸

Try our entire pressure injury prevention portfolio.



ALLEVYN[◊] LIFE



LEAF[◊]
Patient Monitoring System



SECURA[◊] Skin Care
NO-STING SKIN-PREP[◊]
Skin Care

For detailed product information, including indications for use, contraindications, precautions and warnings, please consult the product's applicable Instructions for Use (IFU) prior to use.

*Tested against Mepilex™ Border

REFERENCES: 1. ALLEVYN[◊] LIFE vs Mepilex™ Border – Product Retention Performance Testing. Data on File report HVT080 September 2018 2. Rossington, A. Pressure transmission testing of ALLEVYN LIFE when wet and dry, Mepilex™ Border and Optifoam™ Gentle. Data on File report DS/18/351R- September 2018 3. Earl, K. Wound Model Testing of ALLEVYN LIFE Gen2 using Horse Serum. Data on File report DS/14/303R - April 2016 4. Rossington, A. Product Performance of Next Generation ALLEVYN LIFE. Data on File report GMCA-DOF/08. April 2016. 5. National Pressure Ulcer Advisory Panel, European Pressure Ulcer Advisory Panel and Pan Pacific Pressure Injury Alliance. Prevention and Treatment of Pressure Ulcers: Quick Reference Guide. Emily Haesler (Ed). Cambridge Media: Osborne Park, Australia; 2014 6. Joy H et al. A collaborative project to enhance efficiency through dressing change practice. Journal of Wound Care. Vol 24. No 7. July 2015 P3-4 7. Forni C, D'Alessandro F, Gallerani P, et al. Effectiveness of using a new polyurethane foam multi-layer dressing in the sacral area to prevent the onset of pressure ulcer in the elderly with hip fractures: A pragmatic randomised controlled trial. Int Wound J. 2018; 15(3):383-390. 8. Forni C, Searle R. Economic evaluation of the use of a multi-layer polyurethane foam dressing for the prevention of pressure ulcers in elderly patients with hip fractures. Poster presented at EPUAP annual meeting, September 2018.

Advanced Wound Management

Smith & Nephew, Inc.
Fort Worth, TX 76109
USA

www.smith-nephew.com

Customer Care Center

1-800-876-1261
T 727-392-1261
F 727-392-6914

Ordering codes

Product #	Dressing Sizes	Dressings per box
	Border to border	Pad size
66801306	Sacrum 6 3/4in x 6 7/8in 17.2cm x 17.5cm	4 7/8in x 3 5/8in 8.5cm x 12.4cm
66801307	Sacrum 8 1/2in x 9in 21.6cm x 23cm	6 3/4in x 4 13/16in 12.3cm x 17cm

To order a sample,
visit allevyn.com



Application + Removal Guide

The all-in-one dressing for wound management and pressure injury prevention that stays in place¹, redistributes pressure^{2*} and minimizes leakage.³

* in vitro

Smith+Nephew

ALLEVYN[◊] LIFE
Foam Dressing

Application + Removal Guide for ALLEVYN LIFE Sacrum Foam Dressing

Smith+Nephew

ALLEVYN[◊] LIFE

Foam Dressings

+ Prepare

Cleanse and dry the skin. Ensure it is free of excess hair, dimethicone and other emollients.

To help protect intact or damaged skin, consider using a liquid skin barrier such as NO-STING SKIN-PREP[◊] before applying the dressing.



+ Apply




Select the dressing size to suit the patient, ensuring the dressing pad will cover the area to be protected.




Remove the middle liner first.



Separate the gluteal cleft to ensure complete contact.



Fold with adhesive side down and place narrow end at least 2 centimeters above anal sphincter.




Smooth from midpoint of dressing outward.

+ Inspect for prevention only*



Lift the border from the top of the dressing and peel downward just enough to inspect the skin; leave the dressing intact in the gluteal cleft.



After inspection, smooth the borders back into place, ensuring there are no ridges, folds or air pockets.

Regular skin inspection is important when using the dressing as part of a pressure injury prevention protocol.⁵

Do not roll the edge of the border back onto itself.

+ Remove

When used for wound treatment, change the dressing with each wound assessment. Do not reapply the used dressing.



Lift from the top or proximal edge and peel down slowly toward the anus.

*ALLEVYN LIFE is indicated for pressure injury prevention on intact skin as part of a comprehensive pressure injury prevention protocol.