



Implement repositioning reminder strategies to promote adherence to repositioning regimens
(Guideline Section 5.4)

Use a soft silicone multi-layered foam dressing to protect the skin for individuals at risk of pressure injuries
(Guideline Section 3.5)

Use a prophylactic dressing as an adjunct to heel offloading and other strategies to prevent heel pressure injuries
(Guideline Section 6.4)

Use a prophylactic dressing beneath a medical device to reduce the risk of medical-related pressure injuries
(Guideline Section 8.5)

ALLEVYN GENTLE BORDER demonstrated a relative reduction in Stage I pressure injury incidence (p=0.006) of **90%*30** ↓

ALLEVYN LIFE was shown in one study to reduce sacral pressure injuries by **71%*7** ↓ and in another study to reduce medical device-related pressure injuries from from 27% to **zero*21** ↓

*As compared to standard preventative care alone and as part of a pressure injury prevention protocol

PATIENT TURNING



LEAF Patient Monitoring System



ALLEVYN Foam Dressings



SECURA Skin Care

Shown in a randomized controlled trial to reduce incidence of pressure injuries by **73%*4** ↓

One facility reported that the LEAF System helped save **\$71,500*40** ↓ in one month, in non-reimbursable treatment costs.

Implement an early mobilization program that increases activity and mobility as rapidly as tolerated
(Guideline Section 5.15)

Reposition patients such that optimal offloading of bony prominences and maximum redistribution of pressure is achieved
(Guideline Section 5.5)

Initiate frequent small shifts in body position for critically ill individuals who are too unstable to maintain a regular repositioning schedule and to supplement regular repositioning
(Guideline Section 5.18)

Keep skin clean (avoid use of alkaline soaps and cleansers) and appropriately hydrated
(Guideline Section 3.1)

Cleanse skin promptly after episodes of incontinence
(Guideline Section 3.1)

Protect skin from moisture with a barrier product
(Guideline Section 3.1)